

University of Groningen

## Changing doctor prescribing behaviour, a systematic review

Gill, P; Freemantle, N.; Makela, M.; Vermeuen, K.; Bond, C.; Thorsen, T.; Ryan, G.; Haaijer-Ruskamp, F.M.

*Published in:*  
Pharmacy World & Science

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1999

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Gill, P., Freemantle, N., Makela, M., Vermeuen, K., Bond, C., Thorsen, T., Ryan, G., & Haaijer-Ruskamp, F. M. (1999). Changing doctor prescribing behaviour, a systematic review. *Pharmacy World & Science*, 21, 158-167.

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Changing doctor prescribing behaviour

- P.S. Gill, M. Mäkelä, K.M. Vermeulen, N. Freemantle, G. Ryan, C. Bond, T. Thorsen and F.M. Haaijer-Ruskamp

*Pharm World Sci* 1999;21(4): 158-167.  
© 1999 Kluwer Academic Publishers. Printed in the Netherlands.

**PS Gill** (correspondence): Department of Primary Care and General Practice, University of Birmingham, Birmingham, B15 2TT, UK (e-mail: p.s.gill@bham.ac.uk)  
Funding: none, Conflict of interest: nil

**M Mäkelä:** Services and Quality Stakes, National Research and Development Centre for Welfare and Health, Helsinki, Finland

**KM Vermeulen:** Northern Centre for Health Care Research and Groningen Utrecht Institute for Drug Exploration, University of Groningen, The Netherlands

**N Freemantle:** Centre for Health Economics, University of York, UK

**G Ryan:** Mercer Southern School of Pharmacy, Atlanta, Georgia, USA

**C Bond:** Department of General Practice, Aberdeen University, UK

**T Thorsen:** Danish Institute for Health Services Research and Development, Copenhagen, Denmark

**FM Haaijer-Ruskamp:** Northern Centre for Health Care Research and Groningen Utrecht Institute for Drug Exploration, University of Groningen, The Netherlands

**Keywords** Control Groups, Intervals, Interventions

## Abstract

The aim of this overview was to identify interventions that change doctor prescribing behaviour and to derive conclusions for practice and further research. Relevant studies (indicating prescribing as a behaviour change) were located from a database of studies maintained by the Cochrane Collaboration on Effective Professional Practice. This register is kept up to date by searching the following databases for reports of relevant research: DHSS-DATA; EMBASE; MEDLINE; SIGLE; Resource Database in Continuing Medical Education (1975-1994), along with bibliographies of related topics, hand searching of key journals and personal contact with content area experts. Randomised controlled trials and non-equivalent group designs with pre- and post-intervention measures were included. Outcome measures were those used by the study authors. For each study we determined whether these were positive, negative or inconclusive. Positive studies (+) were those that demonstrated a statistically significant change in the majority of outcomes measured at level of  $p \leq 0.05$  between the intervention and control groups. Negative studies (-) showed a significant change in the opposite direction and inconclusive studies ( $\approx$ ) showed no significant change compared to control or no overall positive findings. We identified 79 eligible studies which described 96 separate interventions to change prescribing behaviour. Of these interventions, 49 (51%, 41%-61%) showed a positive significant change compared to the control group but interpretation of specific interventions is limited due to wide and overlapping confidence intervals.

Accepted December 1998

## Introduction

In recent years, drug expenditure in many health care systems has grown rapidly and varies from 8.4% (USA) to 25.6% (Portugal) [1]. The total costs of drugs prescribed by general practitioners is four times that of drugs prescribed by hospitals. This rise is due to a number of factors and measures have been introduced within the last decade to limit this expenditure, including shift from secondary to primary care [2]. Prescribing is one of the most important health care interventions, however there are frequently gaps between optimal prescribing and that achieved in

practice [3]. Interventions have been targeted at both patients (e.g. increasing prescription charges) and doctors (e.g. feedback of information on drugs and costs). Despite these measures there remain wide variations in prescribing costs and volume within both the primary and secondary sectors.

In the United Kingdom, over two-thirds of general practitioner contacts end with the issuing of a prescription although decisions made by hospital doctors exert considerable indirect influence on general practitioner prescribing [2].

Reviews of improving prescribing in both hospital [4] and ambulatory settings [5] are not new. Systematic reviews have also been published which examine the broad questions of effectiveness of methods to improve professional practice [6,7,8]. However, focused reviews of effectiveness of specific interventions to change doctor prescribing behaviour are lacking.

Our aim was to identify interventions that affect prescribing behaviour and to derive conclusions for practice and further research. This review was intended to provide an overview of interventions that influence prescribing with subsequent specific reviews following.

## Methods

### Data source

Relevant studies (indicating prescribing as a behaviour change) were located from a database of relevant studies maintained by the Cochrane Collaboration on Effective Professional Practice (CCEPP) up to May 1996 [9]. This register is kept up to date by searching the following databases for reports of relevant research: DHSS-DATA; EMBASE; MEDLINE; SIGLE; Resource Database in Continuing Medical Education (1975-1994), along with bibliographies of related topics [8], hand searching of key journals and personal contact with content area experts.

### Selection of studies

The following inclusion criteria were adopted:-

- Study design - randomised controlled trials (RCT) and non-equivalent group designs with pre post measures. In the latter studies, participants are allocated by some means other than chance to treatment or control groups [10]. Studies were included if the intervention(s) was compared with a no intervention control or normal practice group.
- Types of participants - physicians only (includes those undergoing postgraduate training).
- Types of intervention - professional interventions as described by Bero et al [9].

That is distribution of educational materials; conferences or educational meetings; interventions that involve locally deriving consensus recommendations; educational outreach visits that take place in the participants location; the influence of local opinion lead-

**Table 1** Quality criteria for RCTs

<i>Trial</i>	<i>Concealment of allocation</i>	<i>Follow up of professionals</i>	<i>Follow up of patients/ episodes</i>	<i>Blind Assessment</i>	<i>Baseline measurement</i>	<i>Reliable outcome measure</i>	<i>Protection against contamination</i>	<i>Score</i>
RCT - unit of analysis = physician or practice								
Hershey 1988 (78)	✓	✓	±	✓	✓	✓	✓	13
Schechtman 1995 (79)	±	✓	±	✓	✓	✓	±	11
Johnson 1976 (80)	✓	±	✓	±	✓	✓	±	11
Hershey 1986 (81)	±	✓	±	✓	✓	✓	±	11
Meyer 1991 (82)	±	±	✓	✓	✓	✓	±	11
White 1985 (83)	±	±	±	✓	✓	✓	✓	11
Gelbach 1984 (84)	±	×	±	✓	✓	✓	±	9
Landgren 1988 (85)	±	±	✓	✓	✓	✓	✓	12
Putnam 1985 (86)	±	✓	✓	±	±	✓	±	10
Stergachis 1987 (87)	×	±	✓	✓	±	±	×	7
McDonald 1984 (88)	×	±	±	✓	±	✓	±	8
Karuzza 1995 (89)	±	✓	✓	✓	✓	✓	✓	13
Hopkins 1980 (90)	×	✓	✓	✓	±	±	×	8
Cowan 1992 (91)	×	✓	×	✓	±	×	±	6
Tierney 1986 (92)	±	±	±	✓	±	✓	×	8
Lobach 1994 (93)	±	×	×	×	✓	±	±	5
Gullion 1988 (94)	±	✓	×	✓	±	✓	✓	10
Stergachis 1990 (95)	±	✓	✓	✓	✓	✓	±	12
Avorn 1992 (96)	±	✓	×	±	±	±	✓	8
Avorn 1983 (97)	±	✓	✓	✓	✓	✓	✓	13
Bjornson 1990 (98)	±	±	±	✓	±	±	±	8
Denig 1990 (99)	±	✓	±	✓	✓	✓	✓	12
McDonald 1976 (100)	✓	±	±	±	±	±	±	8
Palmer 1985 (101)	±	±	±	±	×	✓	✓	8
Steele 1989 (102)	✓	✓	✓	✓	✓	✓	±	13
McConnell 1982 (103)	±	✓	✓	✓	×	✓	±	10
Maiman 1988 (104)	±	✓	✓	✓	±	±	±	10
McAlister 1986 (105)	±	✓	±	±	±	±	±	8
Ornstein 1991 (106)	±	✓	×	✓	✓	±	±	9
Soumerai 1993 (107)	±	✓	±	×	×	✓	±	7
Berings 1994 (108)	✓	✓	±	±	✓	±	✓	11
Jones 1993 (109)	✓	✓	±	✓	✓	✓	±	12
RCT - unit of analysis = patient or episode of care								
Weingarten 1989 (110)	×	✓	✓	±	±	±	×	7
Johnson 1978 (111)	×	±	✓	✓	✓	±	×	8
Becker 1989 (112)	±	±	×	±	±	±	×	5
Rogers 1982 (113)	±	±	×	✓	±	±	×	6
Kroenke 1990 (114)	±	±	✓	±	✓	±	±	9
Coe 1977 (115)	±	±	✓	✓	✓	✓	×	10
Garrett 1986 (116)	✓	±	±	±	±	×	×	6
Britton 1991 (117)	×	±	±	✓	✓	✓	±	9
Belcher 1990 (118)	✓	±	×	✓	✓	✓	✓	11
White 1989 (119)	±	✓	×	±	±	±	±	7
Rosser 1992 (120)	✓	±	±	✓	✓	✓	×	11
Manheim 1990 (121)	±	±	±	✓	±	✓	×	8
Bryce 1995 (122)	✓	±	✓	✓	✓	±	×	10
Lipton 1992 (123)	×	✓	✓	✓	✓	±	±	10
Anderson 1994 (124)	±	±	±	×	✓	✓	✓	9
Owens 1990 (125)	±	±	±	✓	✓	✓	×	9
Billault 1995 (126)	±	±	✓	±	✓	±	×	8
Rubenstein 1989 (127)	±	✓	✓	±	✓	±	✓	11
Cheney 1987 (128)	±	±	✓	±	±	✓	±	9
Lipton 1994 (129)	✓	✓	✓	✓	✓	✓	×	12
Rodman 1984 (130)	✓	±	±	✓	✓	✓	✓	12
Rosser 1991 (131)	✓	✓	×	✓	✓	±	±	10
White 1987 (132)	±	✓	✓	✓	✓	✓	±	12
Billi 1987 (133)	×	×	×	✓	✓	±	✓	7
Buffington 1991 (134)	±	±	±	✓	×	✓	✓	9
Wilson 1988 (135)	±	✓	✓	✓	✓	✓	✓	13
Katon 1995 (136)	±	±	✓	✓	✓	±	×	9
Browner 1994 (137)	✓	✓	✓	✓	✓	✓	±	13
White 1984 (138)	✓	±	✓	✓	±	✓	×	10
Linn 1980 (139)	✓	✓	✓	±	±	±	✓	11
German 1987 (140)	±	±	✓	±	✓	✓	±	10
Frame 1994 (141)	±	±	±	✓	✓	✓	±	10

✓ = done; ± = not clear; and × = not done and each given a score of ✓ = 2; ± = 1; and × = 0

ers (educational influentials); patient mediated interventions in which information given to or received from patients is intended to influence professional practice; audit and feedback where physicians receive summary information on their practice over time; reminder systems where physicians receive specific reminders at the time of intervention to enhance a

particular strategy or behaviour; marketing, in which physicians are targeted by interventions similar to those used to market commercially specific desired practices. These were either single or in combination (multifaceted interventions).

One of the authors (PG) verified that all studies met the inclusion criteria.

## Data extraction

Eligible studies were assigned to each author and the following categories of data extracted: 1) study design; 2) setting of care; 3) academic status (i.e. teaching or non-teaching); 4) country; 5) clinical problem/area; 6) types of interventions; and 7) outcome measures used by the study authors. For each study we determined whether these were positive, negative or inconclusive. Positive studies (+) were those that demonstrated a statistically significant change in the majority of outcomes measured at level of  $p \leq 0.05$  between the intervention and control groups. Negative studies (-) showed a significant change in the opposite direction and inconclusive studies ( $\approx$ ) showed no significant change compared to control or no overall positive findings. No attempt was made to contact the authors for missing data.

We estimated the proportion of statistically significant studies and approximate 95% confidence intervals using the approach originally described by Clopper and Pearson and cited in Hedges and Olkin [11].

In addition, quality criteria were assessed independently by each author [9]. PG also checked results of these criteria on 40 of the eligible studies.

## Results

One-hundred and forty-five studies were located from the Cochrane Collaboration on Effective Professional Practice Register of which 66 were excluded. Reasons for exclusion were: 1) interrupted time series [12-21]; 2) does not deal exclusively with physician prescribing behaviour or with professional interventions [22-56]; 3) there was no control/normal practice intervention group [57-69]; 4) data were missing to determine outcomes between intervention and control [70-76]; and 5) includes a summary paper of a trial published in another journal [77].

## Quality criteria

For RCTs the following quality criteria were assessed: concealment of allocation; follow up of professionals;

follow up of patients/episodes; blind assessment; baseline measurement; reliable outcome measure; and protection against contamination (Table 1). Quality criteria for non-equivalent group design studies included: baseline measurement; characteristics of studies using second site as control; blind assessment; protection against contamination; follow-up of patients; reliable outcome measures; and follow-up of professionals (Table 2).

Each item was scored as done; not clear or not done [9]. These were then given scores 2, 1 and 0 respectively. Maximum total score for each study was 14 with a mean of 9.65 (sd 2.01) (Tables 1 and 2). Furthermore, there was no difference in the mean quality score before and after the 1990 date of publication (t-test, mean difference -0.44 [-1.35, 0.47]).

## Settings

Sixty-five (82%) of the studies were based in the USA; 3 each in Canada and the UK; 2 each in Australia and Israel; and 1 each in Mexico, Netherlands, Belgium and France (Table 3).

Over a third (31) of the studies took place in outpatient settings; 25 in community settings; 12 in hospitals; and 10 in two settings. It was not possible to characterise the setting in one study [98].

Twenty-seven took place in wholly teaching and 14 in mixed centres; 12 in non-teaching settings; and in 26 studies the teaching status was not clear.

Interventions included in the evaluations included attempts to modify prescribing practices in a variety of clinical areas [78-82 84 86 97 101-102 107 111 114 116-117 121 123 125 127 129 132-133 144 150-151 154]; chronic conditions such as diabetes mellitus [93 100] and asthma [119 122 142]; cardiovascular [83 94 98 105 113 115 126 130 132]; infections [85 95 103-104 146]; preventive care [88-89 91-92 106 110 112 118 120 124 128 131 134-135 137-138 141 147 152 155-156]; psychiatric [96 108 136 140 150]; gastrointestinal [109 109 145 148]; musculo-skeletal [87,143]; surgical conditions [90 139] and problems in childhood [156].

**Table 2** Quality criteria for non-equivalent group design studies

Trial	Baseline measurement	Characteristics of studies using second site as control	Blind assessment	Protection against contamination	Reliable outcome measures	Follow-up of professionals	Follow-up of patients	Score
Non equivalent group design - physician or practice as unit of analysis								
Gorton 1995 (142)	±	×	±	✓	✓	✓	±	9
Petersen 1995 (143)	✓	×	✓	✓	✓	✓	±	11
McDonald 1976 (144)	±	±	✓	✓	✓	✓	✓	12
Raisch 1990 (145)	✓	✓	✓	✓	×	✓	✓	12
Klein 1981 (146)	✓	✓	✓	±	±	±	±	10
Schreiner 1988 (147)	✓	×	±	✓	±	±	±	8
Gutierrez 1994 (148)	✓	✓	✓	±	±	×	±	9
Ray 1986 (149)	×	×	✓	✓	±	✓	±	8
Mason 1993 (150)	±	±	±	✓	±	±	±	8
Non equivalent group design - patient or episode of care as unit of analysis								
Koepsell 1983 (151)	±	✓	✓	✓	✓	×	×	9
Tape 1993 (152)	±	✓	✓	±	✓	±	✓	11
Margolis 1992 (153)	±	±	✓	×	✓	✓	±	9
Fortner 1985 (154)	✓	✓	✓	±	±	±	±	10
Rodney 1983 (155)	±	±	✓	✓	±	±	±	9
Zimmerman 1994 (156)	✓	±	✓	✓	✓	±	✓	12

✓ = done; ± = not clear; and × = not done and each given a score of ✓ = 2; ± = 1; and × = 0

**Table 3** *Studies of effects of interventions on prescribing behaviour of doctors*

Study	Setting	Academic status	Clinical Area	Intervention	Outcome
RCT - unit of analysis = physician or practice					
Hershey 1988 (78)	USA - outpatient	teaching	All clinical areas	Audit and feedback	≈
Schectman 1995 (79)	USA - outpatient	mixed	Antiulcer drugs	Audit and feedback	≈
Johnson 1976 (80)	USA - outpatient	not clear	All clinical areas	Audit and feedback	≈
Hershey 1986 (81)	USA - outpatient	mixed	All clinical areas	Audit and feedback	+
Meyer 1991 (82)	USA - outpatient	teaching	All clinical areas	Audit and feedback	≈
White 1985 (83)	USA - inpatient	mixed	Acute MI	Conferences + Educational materials	+
Gelbach 1984 (84)	USA - community	teaching	All clinical areas	Educational materials + Audit and feedback	+
Landgren 1988 (85)	Australia - inpatient	mixed	Infections	Educational materials + Outreach	+
Putnam 1985 (86)	USA - mixed	not clear	All clinical areas	Educational materials + Outreach + audit and feedback	≈
Stergachis 1987 (87)	USA - mixed	not clear	non-steriodals	outreach	≈
McDonald 1984 (88)	USA - outpatient	teaching	Influenza	Audit and feedback	+
Karuza 1995 (89)	USA - mixed	mixed	influenza	Conferences + Audit and feedback	+
Hopkins 1980 (90)	USA - inpatient	teaching	Surgical emergencies	educational materials	≈
Cowan 1992 (91)	USA - outpatient	teaching	Vaccinations	Audit and feedback	≈
Tierney 1986 (92)	USA - outpatient	teaching	Preventive care	Audit and feedback	+
Lobach 1994 (93)	USA - community	non-teaching	Diabetes mellitus	Audit and feedback	+
Gullion 1988 (94)	USA - mixed	non-teaching	Hypertension	Conferences + Audit and feedback	≈
stergachis 1990 (95)	USA - outpatient	not clear	Upper respiratory infections	Patient mediated	≈
Avorn 1992 (96)	USA - community	not clear	Psychoactive drugs	educational materials + audit and feedback	≈
Avorn 1983 (97)	USA - community	not clear	Variety of clinical areas	i) educational materials ii) educational materials + audit and feedback	i) ≈ ii) ≈
Bjornson 1990 (98)	USA - not clear	not clear	Heart failure	educational materials	≈
Denig 1990 (99)	Netherlands - community	unclear	Irritable Bowel Syndrome	educational materials	≈
McDonald 1976 (100)	USA - outpatient	non-teaching	Diabetes	audit and feedback	+
Palmer 1985 (101)	USA - community	mixed	Variety of clinical areas	educational materials + conferences + audit and feedback	≈
Steele 1989 (102)	USA - outpatient	teaching	Variety of clinical areas	i) Outreach + audit and feedback ii) Audit and feedback	≈ ≈
McConnell 1982 (103)	USA - outpatient	non-teaching	Upper respiratory tract infections	Outreach	+
Maiman 1988 (104)	USA - community	not clear	Otitis media	i) Educational materials ii) Educational materials + conferences	i) + ii) ≈
McAlister 1986 (105)	Canada - community	not clear	Hypertension	Educational materials + patient mediated + audit and feedback	≈
Ornstein 1991 (106)	USA - community	teaching	Preventive care	i) conferences + patient mediated + audit and feedback ii) conferences + audit and feedback iii) conferences + patient mediated	i) ≈ ii) ≈ iii) ≈
Soumerai 1993 (107)	USA - inpatient	mixed	Blood transfusion	conferences + distribution of educational methods + outreach + marketing	+
Berings 1994 (108)	Belgium - outpatient	non-teaching	Anxiety/insomnia	i) educational materials ii) educational materials + outreach	i) + ii) +
Jones 1993 (109)	England - community	mixed	Dyspepsia	educational materials	+
RCT - unit of analysis = patient or episode of care					
Weingarten 1989 (110)	Israel - community	not clear	Preventive care	Audit and feedback	+
Johnson 1978 (111)	USA - outpatient	non-teaching	All clinical areas	Audit and feedback	≈
Becker 1989 (112)	USA - outpatient	teaching	Preventive care	i) Audit and feedback + patient mediated ii) Audit and feedback	i) ≈ ii) ≈
Rogers 1982 (113)	USA - outpatient	teaching	Hypertension	Audit and feedback	≈
Kroenke 1990 (114)	USA - outpatient	teaching	All clinical areas - Elderly	Audit and feedback	≈
Coe 1977 (115)	USA - outpatient	not clear	Hypertension	Audit and feedback	≈
Garrett 1986 (116)	USA - outpatient	not clear	Variety of clinical areas	Audit and feedback	+
Britton 1991 (117)	USA - outpatient	not clear	Variety of Clinical areas	Audit and feedback	≈
Belcher 1990 (118)	USA - outpatient	mixed	Preventive care	Educational materials + conferences + patient reminders + audit and feedback	≈
White 1989 (119)	England - community	not clear	Asthma	Educational materials + patient reminders + audit and feedback	≈
Rosser 1992 (120)	USA - community	teaching	Preventive care	i) Patient mediated - telephone ii) Patient mediated - letter iii) audit and feedback - computer	i) + ii) + iii) +

**Table 3 Continued**

Manheim 1990 (121)	USA - inpatient	teaching	All clinical areas	Educational materials + conferences + audit and feedback	≈
Bryce 1995 (122)	Scotland - community	not clear	Asthma	Educational materials + patient reminders + audit and feedback	+
Lipton 1992 (123)	USA - mixed	non-teaching	All clinical areas	Patient mediated	+
Anderson 1994 (124)	USA - inpatient	mixed	Preventive care	i) Educational materials + outreach	i) +
				ii) Educational materials + outreach + audit and feedback	ii) +
Owens 1990 (125)	USA - inpatient	teaching	All clinical areas	Audit and feedback + patient mediated	+
Billault 1995 (126)	France - mixed	not clear	hypertension	Patient mediated	≈
Rubenstein 1989 (127)	USA - community	non-teaching	various clinical areas	Educational materials + conferences + audit and feedback	≈
Cheney 1987 (128)	USA - outpatient	teaching	Vaccinations	Audit and feedback	+
Lipton 1994 (129)	USA - inpatient	non-teaching	various clinical areas	Patient mediated + audit and feedback	+
Rodman 1984 (130)	USA - inpatient	teaching	CCU	audit and feedback	≈
Rosser 1991 (131)	Canada - community	teaching	Preventive care	i) audit and feedback - computerised decision support	i) +
				ii) Patient mediated - letters	ii) +
				iii) Patient mediated - telephone	iii) +
White 1987 (132)	USA - inpatient	teaching	Warfarin	audit and feedback - computerised decision support	≈
Billi 1987 (133)	USA - inpatient	teaching	various clinical areas	Educational materials + conferences	+
Buffington 1991 (134)	USA - community	not clear	Vaccination	i) audit and feedback	i) +
				ii) audit and feedback + Patient mediated - letter	ii) +
Wilson 1988 (135)	Canada - community	mixed	Smoking cessation	i) Outreach	≈
				ii) Educational materials + conferences	+
Katon 1995 (136)	USA - community	non-teaching	Depression	conferences + Audit and feedback + patient mediated	+
Browner 1994 (137)	USA - outpatient	non-teaching	Hypercholesterolaemia	i) conferences	≈
				ii) conferences + educational materials	≈
White 1984 (138)	USA - inpatient	teaching	Digoxin intoxication	Audit and feedback	+
Linn 1980 (139)	USA - mixed	mixed	Burn care	Educational materials + conferences + audit and feedback	+
German 1987 (140)	USA - outpatient	teaching	Mental health problems	Audit and feedback	≈
Frame 1994 (141)	USA - community	non-teaching	Vaccination	Audit and feedback - computerised decision support	+
Non equivalent group design - physician or practice as unit of analysis					
Gorton 1995 (142)	USA - mixed	mixed	Asthma	i) Educational materials + conferences	i) ≈
				ii) Educational materials + conferences (computer)	ii) ≈
				iii) Educational materials + conferences (multimedia)	iii) ≈
Petersen 1995 (143)	Tasmania - community	not clear	Allopurinol	Educational materials + outreach	+
McDonald 1976 (144)	USA - outpatient	not clear	Various clinical areas	audit and feedback - computerised decision support	+
Raisch 1990 (145)	USA - community	not clear	Antiulcer drugs	Educational materials + outreach	≈
Klein 1981 (146)	USA - outpatient	teaching	Urinary tract infections	conferences	+
Schreiner 1988 (147)	USA - outpatient	teaching	Vaccination	audit and feedback	+
Gutierrez 1994 (148)	Mexico - community	not clear	Diarrhoea	Conferences + audit and feedback	+
Ray 1986 (149)	USA - mixed	not clear	Diazepam	Educational materials + outreach	≈
Mason 1993 (150)	USA - outpatient	teaching	Variety of clinical areas	Outreach	+
Non equivalent group design - patient or episode of care as unit of analysis					
Koepsell 1983 (151)	USA - outpatient	teaching	All clinical areas	Audit and feedback - computerised decision support	≈
Tape 1993 (152)	USA - outpatient	mixed	Vaccination	Audit and feedback	+
Margolis 1992 (153)	Israel - community	not clear	Childhood problems	Audit and feedback - computerised decision support	+
Fortner 1985 (154)	USA - community	not clear	All clinical areas	conferences + patient mediated + audit and feedback	+
Rodney 1983 (155)	USA - outpatient	teaching	Vaccination	Audit and feedback + conferences	+
Zimmerman 1994 (156)	USA - mixed	not clear	Histamine antagonists	Educational materials + audit and feedback	+

+ = significant outcome; ≈ = non-significant or equivocal outcome

Interventions

Within the 79 eligible studies [78-156], there were 96 separate interventions apart from the non-intervention controls.

To aid presentation, the nine interventions detailed above were grouped into six categories: distribution of educational materials; audit and feedback (includes local consensus processes and reminders); outreach (- includes local opinion leaders); patient mediated; conferences; and marketing.

Table 3 shows that 53 studies reported single interventions and the remaining 26 studies were multifaceted. Of the single interventions, audit and feedback were most common.

Forty-nine (49/96) of the interventions showed a positive significant change compared to the control group (Table 4). Note that none of the studies showed a negative effect as we were interested in those studies that demonstrated a statistically significant change in the majority of outcomes.

Studies that have used patient mediated interventions only yield the highest proportion of positive results. Next come interventions which provide outreach ('academic detailing') and audit and feedback (including reminders). The least effective were distribution of educational materials (43%, 13%-78%).

Of the multifaceted interventions, just under fifty percent (49%, 20%-80%) were positive.

Overall, 51% (41%-61%) of the intervention demonstrated a significant change.

The number of positive results was lower in studies

that were set in the USA compared to the other countries (table 5).

There is no difference in the studies reporting a positive result with the proportion of studies above and below the median quality score (table 6).

Discussion

This overview has shown that a number of studies, in primary and secondary settings, are available which address interventions that tend to influence prescribing. The interventions are varied and have been used in various combinations. Further, the majority have been set in the USA and the results may not apply elsewhere.

It is interesting that inspite of the high proportion of studies that show positive results that were unlikely to be accounted for by chance, no clear differences between approaches can be seen, with the exception of the distribution of printed educational materials which has been used widely in combination with other active interventions, but where used alone has proven unimpressive.

Vote counting is an inefficient use of the statistical information in studies, and the poor reporting of results makes the use of more formal methods for pooling data and calculating effect size problematic [11].

Publication bias is the most important potential source of bias in systematic reviews [157]. Although considerable efforts have gone into locating relevant

Table 4 Types of interventions aimed at changing doctor prescribing behaviour

		No. of interventions	Positive findings	% Positive interventions	95% CI
Intervention	distribution of educational materials	7	3	43	13, 78
	audit and feedback	33	17	52	34, 66
	outreach	4	2	50	10, 90
	patient mediated	8	5	63	30, 90
	conferences	1	1	100	-
	marketing	0	0	0	-
	multifaceted*	43	21	49	20, 80
Overall		96	49	51	41, 61

\* two or more of the above interventions

Table 5 Country of study

Country	Positive interventions	No of interventions	% Positive interventions (95% CI)
USA	36	78	46 (33, 59)
other	13	18	72 (43, 95)

Table 6 Outcome of studies against score

		Positive findings	Total no of studies	% Positive studies (95% CI)
Median quality score	≤9	18	37	49 (32, 65)
	≥10	22	42	52 (34, 63)

studies, it is likely that some have not been included in this review. We decided a priori not to include time series designs. Auto regressive integrated moving average models which are widely used in time series analysis [10], are required to take into account the non independent nature of observations from the same population over time, and this is very rarely robustly undertaken in the literature [3]. However, the relatively small number of time series data sets identified would be unlikely to reverse the results from other included designs in this review. Although randomised trials provide the most valid estimates of effect, time series designs deserve further consideration in the future as they may be used in situations where experimental designs are not possible, such as national guideline implementation programmes or mass media interventions. We found 9 non-randomised trials, which are potentially open to bias, and it was not clear in most cases why a decision had been made not to allocate by chance (and thus increase the validity of the trial results). In future trials randomisation should be considered more frequently [158].

### Study limitations

Even though the CCEPP database has been rigorously developed publication bias has to be considered as the search strategy may not have identified important negative results from trials. In addition the database includes studies that use quantitative methodologies (RCTs, Controlled trials and Interrupted Time Series) and valuable results from qualitative studies which provide insight into behaviour change have been excluded.

Data extraction has been difficult, as the reporting of details and results has been incomplete or vague. We did not contact the authors for further clarification of data.

Some of these studies used multiple outcome measures (e.g. number prescriptions/patient and % generic prescribing) and for these studies, a positive finding was that which had majority significant findings. Studies are also subject to another analysis error where, for example, the unit of allocation is a health care provider but the analysis is based upon the number of patients or pills. This could under power the individual study and thereby increase the chance of a spuriously significant finding [159].

### Implications

This review has shown that 51% of interventions studied may have changed prescribing behaviour and a number of research issues are raised. For example, distribution of educational materials alone appear to have little and multifaceted interventions some effect on changing behaviour. Interventions which change doctor prescribing behaviour are complex - there are no "magic bullets" [7]. Further work, in collaboration with the Cochrane Collaboration on Effective Professional Practice [9], is advocated on disentangling the characteristics of these effective interventions that acknowledge the personal, environmental and behavioural factors [160-161]. Also, primary studies are needed which take into account these effective interventions; are multidisciplinary; and which combine both the quantitative and qualitative methodologies.

### Acknowledgement

We are grateful to the CCEPP Editorial Base, York, for the use of the database and in particular to Jo Bark for assistance with this study.

### References

- 1 OECD Health Data 96. A software for the comparative analysis of 27 health systems. CREDES: OECD, 1996.
- 2 Audit Commission. A Prescription for Improvement. Towards more Rational prescribing in General Practice. London: HMSO, 1994.
- 3 Bloor K, Freemantle N. Lessons from international experience in controlling pharmaceutical expenditure II: Influencing doctors. *BMJ* 1996;312:1525-7.
- 4 Soumerai SB, Avorn J. Efficacy and Cost-containment in Hospital Pharmacotherapy: State of the Art and Future Directions. *Milbank Memorial Fund Quarterly/Health and Society* 1984;62:447-74.
- 5 Soumerai SB, McLaughlin TJ, Avorn J. Improving Drug Prescribing in Primary Care: A Critical Analysis of the Experimental Literature. *Milbank Quarterly* 1989;67:268-317.
- 6 Davis DA, Thomson MA, Oxman AD, Haynes RB. Evidence for the Effectiveness of CME: a review of 50 randomised controlled trials. *JAMA* 1992;268:1111-7.
- 7 Oxman AD, Thomson MA, Davis DA, Haynes RB. No magic bullets: a systematic review of 102 trials of interventions to improve professional practice. *Can Med Assoc J* 1995;153:1423-31.
- 8 Effective Health Care Bulletin (1994). Implementing Clinical Practice Guidelines. Bulletin No 8. Leeds: University of Leeds.
- 9 Bero L, Grilli R, Grimshaw J, Oxman AD (eds). In: Cochrane Collaboration on Effective Professional Practice (CCEPP) Module of The Cochrane Database of Systematic Reviews, [updated 02 December 1996]. Available on The Cochrane Library [database on disc and CDROM]. The Cochrane Collaboration; Issue 1. Oxford: Update Software; 1997. Updated quarterly.
- 10 Cook TD, Campbell DT. Quasi-experimentation: design and analysis issues for field settings. Chicago: Rand McNally, 1979.
- 11 Hedges LV, Olkin I. Statistical methods for meta analysis. London: Academic Press, 1985.
- 12 Avorn J, Soumerai SB, Taylor W, et al. Reduction of incorrect antibiotic dosing through a structured educational order form. *Arch Intern Med* 1988;148:1720-4.
- 13 Berbatis CG, Maher MJ, Plumridge RJ, Stoelwinder JU, Zubrick SR. Impact of a drug bulletin on prescribing oral analgesics in a teaching hospital. *Am J Hosp Pharm* 1982;39:98-100.
- 14 Bottiger LE, Westerholm B. Drug-induced blood dyscrasias in Sweden *BMJ* 1973;3:339-43.
- 15 Evans RS, Classen DC, Pestotnik SL, Lundsgaarde, Burke JP. Improving empiric antibiotic selection using computer decision support. *Arch Intern Med* 1994; 154: 878-84.
- 16 Everitt DE, Soumerai SB, Avorn J, Klapholz H, Wessels M. Changing surgical antimicrobial prophylaxis practices through education targeted at senior department leaders. *Infect Control Hosp Epidemiology* 1990; 11(11): 578-83
- 17 Gurwitz JH, Noonan JP, Soumerai SB. Reducing the use of H2-receptor antagonists in the long-term- care setting. *J Am Geriatr Soc* 1992;40(4):359-64.
- 18 Ross-Degnan D, Soumerai SB, Fortess EE, Gurwitz JH. Examining product risk in context: market withdrawal of Zomepirac as a case study. *JAMA* 1993;270(16):1937-42.
- 19 Soumerai SB, Avorn J, Gortmaker S and Hawley S. Effect of government and commercial warnings on reducing prescription misuse: the case of propoxyphene. *Am J Public Health*. 1987;77:1518-23.
- 20 Soumerai SB, McLaughlin TJ, Ross-Degnan D, Casteris CS, Bollini P. Effects of limiting Medicaid drug-reimbursement benefits on the use of psychotropic agents and acute mental health services by patients with schizophrenia. *N Eng J Med* 1994;331:650-5.
- 21 Soumerai SB, Avorn J, Taylor WC, Wessels M, Maher D, Hawley SL. Improving choice of prescribed antibiotics through concurrent reminders in an educational order form. *Med Care* 1993;31(6):552-8.
- 22 Bass MJ, McWhinney IR, Donner A. Do family physicians need medical assistants to detect and manage hypertension? *Can Med Assoc J* 1986;134:1247-55.
- 23 Brown LF, Keily PA, Spencer AJ. Evaluation of a continuing medical education intervention 'periodontics in general practice. *Community Dent Oral Epidemiol* 1994;22:441-7.
- 24 Cohen DI, Littenberg B, Wetzel C, Neuhaser DvB. Improving physician compliance with preventive medicine guidelines. *Med Care* 1982;XX:1040-5.



- 25 Dickinson GR, Holzemer WL, Nichols E. Evaluation of an arthritis continuing education program. *The Journal of Continuing Education in Nursing* 1985;16 (4):127-31.
- 26 Epstein LM, Avni A, Hopp C, Flug D. Evaluation of a program of aftercare for patients discharged from the hospital. *Med Care* 1973;11(4):320-7.
- 27 Fabacher D, Josephson K, Pietruszka F, Linderborn K, Morley JE, Rubenstein LZ. An in-home preventive assessment program for independent older adults: a randomized controlled trial. *J Am Geriatr Soc* 1994;42:630-8.
- 28 Greenfield S, Friedland G, Scifers S, Rhodes A, Black WL, Komaroff AL. Protocol management of dysuria, urinary frequency, and vaginal discharge. *J Fam Practice* 1975; 2(3):179-84.
- 29 Harris BL, Stergachis A, Ried LD. The effect of drug co-payments on utilization and cost of pharmaceuticals in a health maintenance organization. *Med Care* 1990;28(10):907-17.
- 30 Hastings JEF, Mott FD, Barclay A, Hewitt D. Prepaid group practice in Sault Ste. Marie, Ontario: Part I: analysis of utilization records. *Med Care* 1973;11(2):91-103.
- 31 Hedrick SC, Rothman ML, Chapko M, Inui TS, Kelly JR, Ehreth J. Overview and patient recruitment in the adult day health centre evaluation study. *Med Care* 1993;31(9 suppl): S53-S514.
- 32 Hodnett E, Kaufman K, O'Brien-Pallas L, Chipman M. A Strategy to Promote Research-Based Nursing Care: Effects on Childbirth Outcomes. Final Report of Project No.6606-4076-57E funded by the National Health Research and Development Program, Health Canada. 31 March 1994.
- 33 Inui TS, Yourtee EL and Williamson JW. Improved outcomes in hypertension after physician tutorials: a controlled trial. *Ann Intern Med* 1976; 84: 646-51.
- 34 Jin BW, Kim SC, Mori T, Shimao T. The impact of intensified supervisory activities on tuberculosis treatment. *Tubercle and Lung Disease* 1993; 74(4): 267-72.
- 35 Katon W, von Korff M, Lin E, Bush T, Ormel J. Adequacy and duration of antidepressant treatment in primary care. *Med Care* 1992;30(1):67-76.
- 36 Kimberlin CL, Berardo DH, Pendergast JF, McKenzie LC. Effects of an education program for community pharmacists on detecting drug-related problems in elderly patients. *Med Care* 1993;31(5):451-68.
- 37 Kozma CM, Reeder CE, Lingle EW. Expanding Medicaid drug formulary coverage: effects on utilization of related services. *Med Care* 1990;28(10):963-77.
- 38 Krasnik A, Groenewegen PP, Pedersen PA, von Scholten P, Mooney G, Gottschau A, Flierman HA, Damsgaard MT. Changing remuneration systems: effects on activity in general practice. *BMJ* 1990;300:1698-701.
- 39 Logan AG, Milne BJ, Achber C, Campbell WP, Haynes RB. Cost-effectiveness of a worksite hypertension treatment program. *Hypertension* 1981;3(2):211-8.
- 40 Mackie IC, Worthington HV, Hobson P. An investigation into sugar-containing and sugar-free over-the-counter medicines stocked and recommended by pharmacists in the North Western Region of England. *Brit Dent J* 1993; 175(93): 93-8.
- 41 Melin AL, Bygren LO. Efficacy of the rehabilitation of elderly primary health care patients after short-stay hospital treatment. *Med Care* 1992;30(11):1004-15.
- 42 Naughton BJ, Moran MB, Feinglass J, Falconer J, Williams ME. Reducing hospital costs for the geriatric patient admitted from the emergency department: a randomized trial. *JAGS* 1994;42:1045-9.
- 43 Perera DR, LoGerfo JP, Shulenberg E, Ylvisaker JT, Kirz HL. Teaching sigmoidoscopy to primary care physicians: a controlled study of continuing medical education. *J Fam Pract* 1983;16(4):785-8.
- 44 Peterson AM, Kirkwood CF, Hansen LA. Influence of a decision analysis model on selection of drug therapy. *Am J Hosp Pharm* 1991;48:1477-83.
- 45 Rogers JL, Haring OM. The impact of a computerized medical record summary system on incidence and length of hospitalization. *Med Care* 1979;17:618-30.
- 46 Sanazaro PJ, Worth RM. Concurrent quality assurance in hospital care. Report of a study by private initiative in PSRO. *N Eng J Med* 1978;298(21):1171-7.
- 47 Simmer TL, Nerenz DR, Rutt WM, Newcomb CS, Benfer DW. A randomized, controlled trial of an attending staff service in general internal medicine. *Med Care* 1991;29(7 suppl):J531-40.
- 48 Simon JL, Barton Smith D. Change in location of a student health service: a quasi-experimental evaluation of the effects of distance on utilization. *Med Care* 1973;xi(1):59-67.
- 49 Sloan GA, Gordon GS, Cocks, DL. Hospital drug formularies and use of hospital services. *Med Care* 1993;31(10):851-67.
- 50 Smalley WE, Griffin MR, Fought RL, Sullivan L, Ray WA. Effect of a prior-authorization requirement on the use of nonsteroidal anti-inflammatory drugs by Medicaid patients. *New Eng J Med* 1995;332(24):1612-7.
- 51 Soumerai SB, Avorn J, Ross-Degnan D, Gortmaker S. Payment restrictions for prescription drugs under Medicaid. *New Eng J Med* 1987;317:550-6.
- 52 Soumerai SB, Ross-Degnan D, Avorn J, McLaughlin TJ, Choodnovskiy I. Effects of Medicaid drug-payment limits on admission to hospitals and nursing homes. *New Eng J Med* 1991;325(15):1072-7.
- 53 Soumerai SB, Ross-Degnan D, Gortmaker S, Avorn J. Withdrawing payment for nonscientific drug therapy. *AMA* 1990;263(6):831-9.
- 54 The SUPPORT Principal Investigators. A Controlled Trial to Improve Care for Seriously Ill Hospitalized Patients. The Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments (SUPPORT). *JAMA* 1995;274 (20):1591-98.
- 55 Thompson JF, McGhan WF, Ruffalo RL, Cohen DA, Adamcik B, Segal JL. Clinical Pharmacists Prescribing Drug Therapy in a Geriatric Setting: Outcome of a trial. *J Am Geriatr Soc* 1984;32(2):154-9.
- 56 Wing DS, Duff HJ. Evaluation of a therapeutic drug monitoring program for theophylline in a teaching hospital. *Drug Intelligence and Clinical Pharmacy* 1987;21:702-5.
- 57 Brady WJ, Hissa DC, McConnell M, Wones RG. Should physicians perform their own quality assurance audits? *J Gen Intern Med* 1988;3:560-65.
58. Cohen SJ, Christen AG, Katz BP, Drook CA, Davis BJ, Smith DM, Stookey GK. Counselling medical and dental patients about cigarette smoking: the impact of nicotine gum and chart reminders. *Am J Public Health* 1987;77:313-6.
- 59 De Santis G, Harvey KJ, Howard D, Mashford ML, Moulds RFW. Improving the quality of antibiotic prescription patterns in general practice. *Med J Australia* 1994;160:502-5.
- 60 Evans CE, Haynes RB, Birkett NJ, Gilbert JR, Taylor DW, Sackett DL, Johnston ME, Hewson SA. Does a mailed continuing education program improve physician performance? Results of a randomized trial in antihypertensive care. *JAMA* 1986;255(4):501-4.
- 61 Herman CJ, Speroff T, Cebul RD. Improving compliance with immunization in the older adult: results of a randomized cohort study. *JAGS* 1994;42:1154-9.
- 62 Kendrick T, Burns T, Freeling P. Randomized controlled trial of teaching general practitioners to carry out structured assessments of their long term mentally ill patients. *BMJ* 1995;311:93-8.
- 63 Mazzuca SA, Vinicor F, Einterz RM, Tierney WM, Norton JA, Kalasinski LS. Effects of the clinical environment on physicians' response to postgraduate medical education. *Am Educ Res J* 1990;27(3):473-88.
- 64 Pinkerton RE, Tinanoff N, Willms JL, Tapp JT. Resident physician performance in a continuing education format: does newly acquired information improve patient care? *JAMA* 1980;244(19):2183-5.
- 65 Szczepura A, Wilmot J, Davies C, Fletcher J. Effectiveness and cost of different strategies for information feedback in general practice. *Brit J Gen Pract* 1994;44:19-24.
- 66 Tape TG, Givner N, Wigton RS, Seelig CB, Patil K, Campbell JR. Process in ambulatory care: a controlled clinical trial of computerized records. *SCAMC* 1988;749-52.
- 67 Turner RC, Peden JG, O'Brien K. Patient-carried card prompts vs computer-generated prompts to remind private practice physicians to perform health maintenance measures. *Arch Int Med* 1994;154:1957-60.
- 68 Turner RC, Waivers LE, O'Brien K. The effect of patient-carried reminder cards on the performance of health maintenance measures. *Arch Intern Med* 1990;150:645-7.
- 69 Wing D, Duff HJ. Impact of a therapeutic drug monitoring program for digoxin. *Arch Intern Med* 1987;147(8):1405-8.
- 70 Diwan VK, Wahlström R, Tomson G, Beermann B, Sterky G, Eriksson B. Effects of "group detailing" on the prescribing of lipid-lowering drugs: a randomized controlled trial in Swedish Primary Care. *J Clin Epidemiol* 1995;48:705-11.
- 71 North of England Study of Standards and Performance in General Practice. Medical audit in general practice: effects on doctors' clinical behaviour and the health of patients with common childhood conditions. *BMJ* 1992;304:1480-8.
- 72 Rokstad K, Straand J, Fugelli P. Can drug treatment be improved by feedback on prescribing profiles combined with therapeutic recommendations? A prospective controlled trial in general practice. *J Clin Epidemiol* 1995;48(8):1061-8.
- 73 Schaffner W, Ray WA, Federspiel CF, O'Miller WO. Improving antibiotic practice in office prescribing: a controlled trial of three educational methods. *JAMA* 1983;250:1728-32.
- 74 Stross JK, Bole GG. Evaluation of a continuing education program in rheumatoid arthritis. *Arthritis Rheumatism* 1980;23:846-9.
- 75 Stross JK, Bole GG. Evaluation of an educational program for primary care practitioners on the management of osteoarthritis. *Arthritis Rheumatism* 1985;28:108-11.

- 76 Stross JK, Hiss RG, Watts CM, Davis WK, MacDonald R. Continuing education in pulmonary disease for primary care physicians. *Am Rev Resp Dis* 1983;27:739-46.
- 77 Hawkins DW. Clinical pharmacy functions in ambulatory patient care. *J Clin Pharmacol* 1981;21(5-6):245-50.
- 78 Hershey CO, Goldberg HI, Cohen DI. The effect of computerized feedback coupled with a newsletter upon outpatient prescribing charges. *Med Care* 1988;26(1):88-93.
- 79 Schectman JM, Kanwal NJ, Schroth WS, Elinsky EG. The effect of an education and feedback intervention on group-model and network model health maintenance organization physician prescribing behavior. *Med Care* 1995;33:139-44.
- 80 Johnson RE, Campbell WH, Azevedo DJ, Christensen DB. Studying the impact of patient drug profiles in an HMO. *Med Care* 1976;14(10):799-807.
- 81 Hershey CO, Porter DK, Breslau D, Cohen DI. Influence of simple computerized feedback on prescription charges in an ambulatory clinic: a randomized clinical trial. *Med Care* 1986;24:472-81.
- 82 Meyer TJ, Van Kooten D, Marsh S, Prochazka AV. Reduction of polypharmacy by feedback to clinicians. *J Gen Intern Med* 1991;6:133-6.
- 83 White CW, Albanese MA, Brown DD, Caplan RM. The effectiveness of continuing medical education in changing the behaviour of physicians caring for patients with acute myocardial infarction. *Ann Intern Med* 1985;102:686-92.
- 84 Gehlbach SH, Wilkinson WE, Hammond WE, Clapp NE, Finn AL, Taylor WJ, Rodell M. Improving drug prescribing in a primary care practice. *Med Care* 1984;22:193-201.
- 85 Landgren FT, Harvey KJ, Mashford ML, Moulds RF, Guthrie B, Hemming M. Changing antibiotic prescribing by educational marketing. *Med J Australia* 1988;149:595-9.
- 86 Putnam RW, Curry L. Impact of patient care appraisal on physician behaviour in the office setting. *Can Med J* 1985;132:1025-9.
- 87 Stergachis A, Fors M, Wagner EH, Sims DD, Penna P. Effect of clinical pharmacists on drug prescribing in a primary-care clinic. *Am J Hosp Pharm* 1987;44:525-9.
- 88 McDonald CJ, Hui SL, Smith DM, Tierney WM, Cohen SJ, Weinberger M, et al. Reminders to physicians from an introspective computer medical record: a two year randomized trial. *Ann Intern Med* 1984;100:130-8.
- 89 Karuza J, Calkins E, Feather J, Hershey CO, Katz L, Majeroni B. Enhancing physician adoption of practice guidelines. Dissemination of influenza vaccination guideline using a small-group consensus process. *Arch Intern Med* 1995;155(6):625-32.
- 90 Hopkins JA, Shoemaker WC, Greenfield S, Chang PC, McAuliffe T, Spout RW. Treatment of surgical emergencies with and without an algorithm. *Arch Surg* 1980;115:745-50.
- 91 Cowan JA, Heckerling PS, Parker JB. Effect of a fact sheet reminder on performance of the periodic health examination: A randomized controlled trial. *Am J Prev Med* 1992;8:104-9.
- 92 Tierney WM, Hui SL, McDonald CJ. Delayed feedback of physician performance versus immediate reminders to perform preventive care: effects on physician compliance. *Med Care* 1986;24(8):659-66.
- 93 Lobach DF, Hammond WE. Development and evaluation of a computer-assisted management protocol (CAMP): improved compliance with care guidelines for diabetes mellitus. Proceedings - The annual symposium on computer applications in medical care 1994;787-91.
- 94 Gullion DS, Tschann JM, Adamson TE, Coates TJ. Management of hypertension in private practices: a randomized controlled trial in continuing medical education. *J Cont Educ Health Prof* 1988;8:239-55.
- 95 Stergachis A, Newmann WE, Williams KJ, Schnell MM. The effect of a self-care minimal intervention for colds and glue on the use of medical services. *J Gen Intern Med* 1990;5:23-8.
- 96 Avorn J, Soumerai SB, Everitt DE, Ross-Degnan D, Beers MH, Sherman D, Salem-Schatz SR, Fields D. A randomized trial of a program to reduce the use of psychoactive drugs in nursing homes. *N Engl J Med* 1992;327:168-73.
- 97 Avorn J, Soumerai SB. Improving drug-therapy decisions through educational outreach. *N Engl J Med* 1983;302:1457-63.
- 98 Bjornson DC, Rector TS, Daniels CE, Wertheimer AI, Snowdon DA, Litman TJ. Impact of drug-use review program intervention on prescribing after publication of a randomized clinical trial. *Am J Hosp Pharm* 1990;47:1541-6.
- 99 Denig P, Haaijer-Ruskamp FM, Zijlsling DH. Impact of a drug bulletin on the knowledge, perception of drug utility, and prescribing behavior of physicians. *Ann Pharmacotherapy* 1990;24:87-92.
- 100 McDonald CJ. Use of a computer to detect and respond to clinical events: Its effect on clinician behaviour. *Ann Intern Med* 1976;84:162-7.
- 101 Palmer RH, Louis TA, HSu LN, Peterson HF, Rothrock JK, Strain R, Thompson MS, Wright EA. A randomized controlled trial of quality assurance in sixteen ambulatory care practices. *Med Care* 1985;23(6):751-70.
- 102 Steele MA, Bess DT, Franse VL, Graber SE. Cost effectiveness of two interventions for reducing outpatient prescribing costs. *DICP Annals of Pharmacotherapy* 1989;23:497-500.
- 103 McConnell TS, Cushing AH, Bankhurst AD, Healy JL, McIlvenna PA, Skipper BJ. Physician behaviour modification using claims data: Tetracycline for upper respiratory infection. *West J Med* 1982;137:448-50.
- 104 Maiman LA, Becker MH, Liptak GS, Nazarian LF, Rounds KA. Improving pediatricians' compliance-enhancing practices: a randomized trial. *Am J Dis Child* 1988;142:773-9.
- 105 McAlister NH, Covey HD, Tong C, Lee A, Wigle ED. Randomised controlled trial of computer assisted management of hypertension in primary care. *BMJ* 1986;293:670-4.
- 106 Ornstein SM, Garr DR, Jenkins RC, Rust PF, Aron A. Computer-generated physician and patient reminders. Tools to improve population adherence to selected preventive services. *J Fam Pract* 1991;32:82-90.
- 107 Soumerai SB, Salem-Schatz S, Avorn J, Casteris CS, Ross-Degnan D, Popovsky MA. A controlled trial of educational outreach to improve blood transfusion practice. *JAMA* 1993;270:961-6.
- 108 Berings D, Blondeel I, Habraken H. The effect of industry-independent drug information on the prescribing of benzodiazepines in general practice. *Eur J Clin Pharmacol* 1994;46:501-5.
- 109 Jones RH, Lydeard S, Dunleavy J. Problems with implementing guidelines: a randomised controlled trial of consensus management of dyspepsia. *Qual Health Care* 1993;2:217-21.
- 110 Weingarten MA, Bazel D, Shannon HS. Computerized protocol for preventive medicine: a controlled self-audit in family practice. *Fam Pract* 1989;6(2):120-4.
- 111 Johnson RE, Azevedo DJ, Campbell WH, Christensen DB. Examining physicians' drug order recording behaviour. *Med Care* 1978;16(5):408-16.
- 112 Becker DM, Gomez EB, Kaiser DL, Yoshihisi A, Hodge RH. Improving preventive care at a medical clinic: how can the patient help? *Am J Prev Med* 1989;5(6):353-9.
- 113 Rogers JL, Haring OM, Wortman PM, Watson RA, Goetz JP. Medical information systems: assessing impact in the areas of hypertension, obesity and renal disease. *Med Care* 1982;20(1):63-74.
- 114 Kroenke K, Pinholt EM. Reducing polypharmacy in the elderly. *J Am Geriatr Soc* 1990;38(1):31-6.
- 115 Coe FL, Norton E, Oparil S, Tatar A, Pullman TD. Treatment of hypertension by computer and physician - a prospective controlled study. *J Chronic Dis* 1977;30:81-92.
- 116 Garrett LE Jr, Hammond WE, Stead WW. The effects of computerized medical records on provider efficiency and quality of care. *Meth Inform Med* 1986;25:151-7.
- 117 Britton ML, Lurvey PL. Impact of medication profile review on prescribing in a general medicine clinic. *Am J Hosp Pharm* 1991;48:265-70.
- 118 Belcher DW. Implementing preventive services: success and failure in an outpatient trial. *Arch Intern Med* 1990;150:2533-41.
- 119 White PT, Pharoah CA, Anderson HR, Freeling P. Randomized controlled trial of small group education on the outcome of chronic asthma in general practice. *J R Coll Gen Pract* 1989;39(322):182-6.
- 120 Rosser WW, Hutchison BG, McDowell I, Newell C. Use of reminders to increase compliance with tetanus booster vaccination. *Can Med Assoc J* 1992;146:911-7.
- 121 Manheim LM, Feinglass J, Hughes R, Martin GJ, Conrad K, Hughes EFX. Training house officers to be cost conscious: effects of an educational intervention on charges and length of stay. *Med Care* 1990;28:29-42.
- 122 Bryce FP, Neville RG, Crombie RG, Clark RA, McKenzie P. Controlled trial of an audit facilitator in diagnosis and treatment of childhood asthma in general practice. *BMJ* 1995;310:838-42.
- 123 Lipton HL, Bero LA, Bird JA, McPhee SJ. The impact of clinical pharmacists' consultations on physicians geriatric drug prescribing. *Med Care* 1992;30:646-58.
- 124 Anderson FA, Wheeler HB, Goldberg RJ, Hosmer DW, Forcier A, Patwardhan NA. Changing clinical practice. Prospective study of the impact of continuing medical education and quality assurance programs on use of prophylaxis for venous thromboembolism. *Arch Intern Med* 1994;154:669-677.
- 125 Owens NJ, Sherburne NJ, Silliman RA, Fretwell MD. The Senior Care Study. The Optimal Use of Medications in Acutely Ill Older Patients. *J Am Geriatr Soc* 1990;38(10):1082-7.
- 126 Billault B, Degoulet P, Devries C, Plouin PF, Chatellier G, Menard J. Use of a standardized personal medical record by patients with hypertension: a randomized controlled trial. *MD Computing* 1995;1:31-5.

- 127 Rubenstein LV, Calkins DR, Young RT, Cleary PD, Fink A, Koseoff J, et al. Improving patient function: a randomized trial of functional disability screening. *Ann Intern Med* 1989;111(10):836-42.
- 128 Cheney C, Ramsdell JW. Effect of medical records' checklists on implementation of periodic health measures. *Am J Med* 1987;83:129-36.
- 129 Lipton HL, Bird JA. The Impact of Clinical Pharmacists' Consultations on Geriatric Patients' Compliance and Medical Care Use: A Randomized Controlled Trial. *The Gerontologist* 1994;34(3):307-15.
- 130 Rodman JH, Jelliffe RW, Kolb E, Tuey DB, Guzman MF de, Wagers PW et al. Clinical studies with computer-assisted initial lidocaine therapy. *Arch Intern Med* 1984;703-9.
- 131 Rosser WW, McDowell I, Newell C. Use of reminders for preventive procedures in family medicine. *Can Med Assoc J* 1991;145:807-14.
- 132 White RH, Hong R, Venook AP, Daschbach MM, Murray W, Mungall DR et al. Initiation of warfarin therapy. *J Gen Intern Med* 1987;2:141-8.
- 133 Billi JE, Hejna GF, Wolf FM, Shapiro LR, Stross JK. The effects of a cost-education program on hospital charges. *J Gen Intern Med* 1987;2:306-11.
- 134 Buffington J, Bell KM, LaForce FM and the Genesee Hospital Medical Staff. A target-based model for increasing influenza immunizations in private practice. *J Gen Intern Med* 1991;6:204-9.
- 135 Wilson DM, Taylor DW, Gilbert JR, Best A, Lindsay EA, Willms DG et al. A randomized trial of a family physician intervention for smoking cessation. *JAMA* 1988;260(11):1570-5.
- 136 Katon W, Von Korff M, Lin E, Walker E, Simon GE, Bush T et al. Collaborative Management to Achieve Treatment Guidelines. Impact on Depression in Primary Care. *JAMA* 1995;273(13):1026-31.
- 137 Browner WS, Baron RB, Solkowitz S, Adler LJ, Gullion DS. Physician management of hypercholesterolemia: a randomized trial of continuing medical education. *West J Med* 1994;161:572-8.
- 138 White KS, Lindsay A, Pryor A, Brown WF, Walsh K. Application of a computerized medical decision-making process to the problem of digoxin intoxication. *JACC* 1984;4(3):571-6.
- 139 Linn BS. Continuing medical education. Impact on emergency room burn care. *JAMA* 1980;244(6):565-70.
- 140 German PS, Shapiro S, Skinner EA, von Korff M, Klein LE, Turner RW et al. Detection and management of mental health problems of older patients by primary care providers. *JAMA* 1987;257(4):489-493.
- 141 Frame PS, Zimmer JG, Werth PL, Hall J, Eberly SW. Computer-based vs manual health maintenance tracking: a controlled trial. *Arch Fam Med* 1994;3:581-8.
- 142 Gorton TA, Cranford CO, Golden WE, Walls RC, Pawelak JE. Primary care physicians' response to dissemination of practice guidelines. *Arch Fam Med* 1995; 4(2): 135-42.
- 143 Peterson GM, Sugden JE. Educational program to improve the dosage prescribing of allopurinol. *Med J Australia* 1995;162:74-7.
- 144 McDonald CJ. Protocol-based computer reminders, the quality of care and the non-perfectability of man. *N Engl J Med* 1976;295(24):1351-5.
- 145 Raisch DW, Bootman JL, Larson LN, McGhan WF. Improving antiulcer agent prescribing in a health maintenance organization. *Am J Hosp Pharm* 1990;47:1766-73.
- 146 Klein LE, Charache P and Johannes RS. Effect of physician tutorials on prescribing patterns of graduate physicians. *J Med Educ* 1981;56:504-11.
- 147 Schreiner DT, Petrusa ER, Rettie CS, Kluge RM. Improving compliance with preventive medicine procedures in a house staff training program. *South Med J* 1988;81:1553-7.
- 148 Gutierrez G, Guisafre J, Bronfman M, Walsh J, Martinez H, Munoz O. Changing physician prescribing patterns: evaluation of an educational strategy for acute diarrhoea in Mexico City. *Med Care* 1994;32(5):436-46.
- 149 Ray WA, Blazer DG, Schaffner W, Federspiel CF, Fink R. Reducing long-term diazepam prescribing in office practice: a controlled trial of educational visits. *JAMA* 1986;256(18):2536-9.
- 150 Mason JD, Colley CA. Effectiveness of an ambulatory care clinical pharmacist: a controlled trial. *Ann Pharmacotherapy* 1993;27:555-9.
- 151 Koepsell TD, Gurtel A, Diehr PH, Temkin NR, Helfand KH, Gleser MA et al. The Seattle evaluation of computerized drug profiles: effects on prescribing practices and resource use. *Am J Public Health* 1983;73:850-55.
- 152 Tape TG, Campbell JR. Computerized medical records and preventive health care: success depends on many factors. *Am J Med* 1993;94:619-25.
- 153 Margolis CZ, Warshawsky SS, Goldman L, Dagan O, Wirtshafter D, Pliskin JS. Computerized algorithms and pediatricians' management of common problems in a community clinic. *Acad Med* 1992;67:282-4.
- 154 Fortner CR, Tarrant J and Felton HF. Report on an ambulatory clinical pharmacy program and its cost effectiveness. *Group Health Journal* 1985;6:3-6.
- 155 Rodney WM, Chopivsky P, Quan M. Adult immunization: the medical record design as a facilitator for physician compliance. *J M Educ* 1983;58:576-80.
- 156 Zimmerman DR, Collins TM, Lipowski EE, Sainfort F. Evaluation of a DUR intervention: a case study of histamine antagonists. *Inquiry* 1994;31:89-101.
- 157 Beggs CB, Berlin JA. Publication bias: a problem in interpreting medical data. *J R Statist Soc* 1988;151:419-63.
- 158 Freemantle N, Wood J, Crawford F. Evidence into practice, experimentation and quasi experimentation: are the methods up to the task? *J Epid and Comm Hlth* 1998;52:75-81.
- 159 Campbell MK, Grimshaw JM. Cluster randomised trials: time for improvement. *BMJ* 1998;317:1171-3.
- 160 Bandura A. *Social Foundations of Thought and Action: A Social Cognitive Theory*. Englewood Cliffs, NJ: Prentice Hall, 1986.
- 161 Armstrong D, Reyburn H, Jones R. A study of general practitioner's reasons for changing their prescribing behaviour. *BMJ* 1996;949-52.